arXiv:0704.2858v5 [math.AG] 13 May 2007

STUDIES ON THE CALOGERO-MOSER SYSTEM OF TYPE A_1

YUSUKE SASANO

Abstract. We study the Calogero-Moser system of type A_1.

0. Rational model

The Calogero-Moser system of rational model is given by

$$\frac{dq_1}{dt}=\frac{\partial H_{rational}}{\partial p_1},\quad \frac{dp_1}{dt}=-\frac{\partial H_{rational}}{\partial q_1},\quad \frac{dq_2}{dt}=\frac{\partial H_{rational}}{\partial p_2},\quad \frac{dp_2}{dt}=-\frac{\partial H_{rational}}{\partial q_2}, \tag{1}$$

$$H_{rational}(q_1,p_1,q_2,p_2,t;g)=\frac{p_1^2+p_2^2}{2}+\frac{g^2}{2(q_1-q_2)^2}.$$

The system (1) has the following symmetry:

$$\pi :(q_1,p_1,q_2,p_2,t;g) \rightarrow (q_2,p_2,q_1,p_1,t;g). \tag{2}$$

In this paper, we present a polynomial Hamiltonian $H_{polynomial}$ by an explicit birational and symplectic transformation φ.

Theorem 0.1. *By the following birational and symplectic transformation*

$$\varphi : (q_1,p_1,q_2,p_2) \rightarrow \left(1/(q_1-q_2),-(q_1-q_2)^2p_1,q_2,p_2+p_1\right), \tag{3}$$

we can obtain the polynomial Hamiltonian system:

$$\begin{cases} \dfrac{dq_1}{dt}=p_2q_1^2+2q_1^4p_1,\\ \dfrac{dp_1}{dt}=-4q_1^3p_1^2-2q_1p_1p_2-g^2q_1,\\ \dfrac{dq_2}{dt}=p_2+q_1^2p_1,\\ \dfrac{dp_2}{dt}=0 \end{cases} \tag{4}$$

with the polynomial Hamiltonian $H_{polynomial}$

$$H_{polynomial}(q_1,p_1,q_2,p_2,t;g)=q_1^2p_1p_2+q_1^4p_1^2+\frac{g^2q_1^2}{2}+\frac{p_2^2}{2}. \tag{5}$$

Theorem 0.2. *The system* (4) *has two first integrals*

$$I_1:=H_{polynomial},\ \ I_2:=p_2 \tag{6}$$

satisfying the relation

$$\{I_1,I_2\}=0, \tag{7}$$

where $\{,\}$ is the Poisson bracket such that $\{q_i,p_j\}=\delta_{ij}$.

2000 Mathematics Subject Classification: 34M55,34M45,58F05,32S65.

Theorem 0.3. *By eliminating p_1, the Hamiltonian system*

$$\begin{cases} \dfrac{dq_1}{dt} = \dfrac{\partial K}{\partial p_1} = cq_1^2 + 2q_1^4 p_1, \\ \dfrac{dp_1}{dt} = -\dfrac{\partial K}{\partial q_1} = -4q_1^3 p_1^2 - 2cq_1 p_1 - g^2 q_1 \quad (c \in \mathbb{C}), \\ (K = cq_1^2 p_1 + q_1^4 p_1^2 + \dfrac{g^2 q_1^2}{2}) \end{cases} \tag{8}$$

is equivalent to the equation for $y := q_1$

$$\begin{aligned} \frac{d^2 y}{dt^2} =& -2g^2 y^5 + \frac{2}{y}\left(\frac{dy}{dt}\right)^2 \\ =& \frac{2\left(\frac{dy}{dt} - gy^3\right)\left(\frac{dy}{dt} + gy^3\right)}{y}. \end{aligned} \tag{9}$$

For (8) we easily see that

$$p_1 = \frac{\frac{dq_1}{dt} - cq_1^2}{2q_1^4}. \tag{10}$$

By using (8) and (10), we can obtain (9).

We also find some Bäcklund transformations of the system (4).

Theorem 0.4. *The system (4) becomes again a polynomial Hamiltonian system by each symplectic transformation:*

$$\begin{aligned} r_1 :& x_1 = -q_1 p_1^2, \ y_1 = \frac{1}{p_1}, \ z_1 = q_2, \ w_1 = p_2, \\ r_2 :& x_2 = q_1 q_2, \ y_2 = \frac{p_1}{q_2}, \ z_2 = \frac{1}{q_2}, \ w_2 = -(q_2 p_2 - q_1 p_1) q_2, \\ r_3 :& x_3 = -(q_1 p_1 - q_2 p_2) p_1, \ y_3 = \frac{1}{p_1}, \ z_3 = \frac{q_2}{p_1}, \ w_3 = p_2 p_1, \\ r_4 :& x_4 = \frac{q_1}{p_2}, \ y_4 = p_1 p_2, \ z_4 = q_2 - \frac{q_1 p_1}{p_2}, \ w_4 = p_2, \\ r_5 :& x_5 = q_1, \ y_5 = p_1 - \frac{q_2 p_2}{q_1}, \ z_5 = q_2 q_1, \ w_5 = \frac{p_2}{q_1}, \\ r_6 :& x_6 = q_1, \ y_6 = p_1 + \frac{c}{q_1}, \ z_6 = q_2, \ w_6 = p_2 \quad (c \in \mathbb{C}), \\ r_7 :& x_7 = q_1, \ y_7 = p_1, \ z_7 = q_2 + \frac{c}{p_2}, \ w_7 = p_2 \quad (c \in \mathbb{C}), \\ r_8 :& x_8 = q_1, \ y_8 = p_1, \ z_8 = \frac{1}{q_2}, \ w_7 = -q_2^2 p_2. \end{aligned} \tag{11}$$

Corollary 0.1. *The system (4) passes the Painlevé test.*

1. Hyperbolic model

The Calogero-Moser system of hyperbolic model is given by

$$\begin{aligned}&\frac{dq_1}{dt}=\frac{\partial H_{rational}}{\partial p_1}, \quad \frac{dp_1}{dt}=-\frac{\partial H_{rational}}{\partial q_1}, \quad \frac{dq_2}{dt}=\frac{\partial H_{rational}}{\partial p_2}, \quad \frac{dp_2}{dt}=-\frac{\partial H_{rational}}{\partial q_2},\\ &H_{rational}(q_1,p_1,q_2,p_2,t;g)=\frac{p_1^2+p_2^2}{2}-\frac{2g(g-1)q_1q_2}{(q_1-q_2)^2}.\end{aligned} \tag{12}$$

The system (12) has the following symmetry:

$$s:(q_1,p_1,q_2,p_2,t;g)\rightarrow(q_2,p_2,q_1,p_1,t;g). \tag{13}$$

In this paper, we present a polynomial Hamiltonian $H_{polynomial}$ by an explicit birational and symplectic transformation φ.

Theorem 1.1. *By the following birational and symplectic transformation*

$$\varphi:(q_1,p_1,q_2,p_2)\rightarrow\left(1/(q_1-q_2),-(q_1-q_2)^2p_1,q_2,p_2+p_1\right), \tag{14}$$

we can obtain the polynomial Hamiltonian system:

$$\begin{cases}\dfrac{dq_1}{dt}=q_1^2(2q_1^2p_1+p_2),\\ \dfrac{dp_1}{dt}=-2(q_1p_1p_2+2q_1^3p_1^2-g(g-1)q_2(2q_1q_2+1),\\ \dfrac{dq_2}{dt}=q_1^2p_1+p_2,\\ \dfrac{dp_2}{dt}=2g(g-1)q_1(2q_1q_2+1)\end{cases} \tag{15}$$

with the polynomial Hamiltonian $H_{polynomial}$

$$H_{polynomial}(q_1,p_1,q_2,p_2,t;g)=\frac{p_2^2+2q_1^2p_1p_2+2q_1^4p_1^2-4g(g-1)q_1q_2(q_1q_2+1)}{2}. \tag{16}$$

We remark that the system (15) has (16) as its first integral.

Proposition 1.1. *The system* (15) *has a 3-parameter family of polynomial solutions*:

$$(q_1,p_1,q_2,p_2)=(0,2g(g-1)(\frac{\alpha_1t^2}{2}+\alpha_2t)+\alpha_3,\alpha_1t+\alpha_2,\alpha_1). \tag{17}$$

Here α_i $(i=1,2,3)$ are constant parameters.

Theorem 1.2. *By eliminating p_1, p_2, the Hamiltonian system* (15) *is equivalent to the equations for $y:=q_1,\ w:=q_2$*

$$\begin{cases}\dfrac{d^2y}{dt^2}=\dfrac{2}{y}\left(\dfrac{dy}{dt}\right)^2+2g(g-1)y^3(2yw+1)^2,\\ \dfrac{d^2w}{dt^2}=2g(g-1)y(yw+1)(2yw+1).\end{cases} \tag{18}$$

We also find some Bäcklund transformations of the system (15).

Theorem 1.3. *The system* (15) *becomes again a polynomial Hamiltonian system by each symplectic transformation*:

$$
\begin{aligned}
&r_1 : x_1 = -q_1 p_1^2, \ y_1 = \frac{1}{p_1}, \ z_1 = q_2, \ w_1 = p_2,\\
&r_2 : x_2 = q_1 q_2, \ y_2 = \frac{p_1}{q_2}, \ z_2 = \frac{1}{q_2}, \ w_2 = -(q_2 p_2 - q_1 p_1) q_2,\\
&r_3 : x_3 = -(q_1 p_1 - q_2 p_2) p_1, \ y_3 = \frac{1}{p_1}, \ z_3 = \frac{q_2}{p_1}, \ w_3 = p_2 p_1,\\
&r_4 : x_4 = \frac{q_1}{p_2}, \ y_4 = p_1 p_2, \ z_4 = q_2 - \frac{q_1 p_1}{p_2}, \ w_4 = p_2,\\
&r_5 : x_5 = q_1, \ y_5 = p_1 - \frac{q_2 p_2}{q_1}, \ z_5 = q_2 q_1, \ w_5 = \frac{p_2}{q_1},\\
&r_6 : x_6 = q_1, \ y_6 = p_1 + \frac{c}{q_1}, \ z_6 = q_2, \ w_6 = p_2 \quad (c \in \mathbb{C}).
\end{aligned} \tag{19}
$$

Corollary 1.1. *The system* (15) *passes the Painlevé test.*

Takasaki discovered a two-parameter family of coupled Painlevé II systems in dimension four with a rational Hamiltonian $H_{\text{rational}} \in \mathbb{C}(q_1, p_1, q_2, p_2, t)$

(20)
$$
\frac{dq_1}{dt} = \frac{\partial H_{\text{rational}}}{\partial p_1} \quad \frac{dp_1}{dt} = -\frac{\partial H_{\text{rational}}}{\partial q_1} \quad \frac{dq_2}{dt} = \frac{\partial H_{\text{rational}}}{\partial p_2} \quad \frac{dp_2}{dt} = -\frac{\partial H_{\text{rational}}}{\partial q_2}
$$

with the Hamiltonian

$$
H_{\text{rational}} = H_{II}(q_1, p_1, t; \alpha_1) + H_{II}(q_2, p_2, t; \beta_1) + \frac{g_4^2}{(q_1 - q_2)^2}. \tag{21}
$$

Here the symbol $H_{II}(q, p, t; \alpha)$ denotes a Hamiltonian of the Painlevé II system:

$$
H_{II}(q, p, t; \alpha_1) = \frac{p^2}{2} - \left(q^2 + \frac{t}{2}\right) p - \alpha_1 q \quad (\alpha_0 + \alpha_1 = 1). \tag{22}
$$

Proposition 1.2. *The Hamiltonian system with*

$$
\tilde{H}_{\text{rational}} = \tilde{H}_{II}(q_1, p_1, t; \alpha_0, \alpha_1) + \tilde{H}_{II}(q_2, p_2, t; \beta_0, \beta_1) + \frac{g_4^2 q_1^2 q_2^2}{(q_1 - q_2)^2} \tag{23}
$$

is equivalent to the system (20) *by the symplectic transformation*

$$
\phi : (q_1, p_1, q_2, p_2) \rightarrow (\frac{1}{q_1}, -(q_1 p_1 + \alpha_1) q_1, \frac{1}{q_2}, -(q_2 p_2 + \beta_1) q_2).
$$

Here the symbol $\tilde{H}_{II}(q, p, t; \alpha)$ denotes a Hamiltonian of the Painlevé II system:

$$
\tilde{H}_{II}(q, p, t; \alpha) = \frac{q^4 p^2}{2} + \alpha q^3 p + \frac{t}{2} q^2 p + \frac{\alpha^2 q^2}{2} + \frac{\alpha t}{2} q + p. \tag{24}
$$

Theorem 1.4. *The system* (20) *is invariant under the following transformations* s_1, s_2 :

(25)
$$
\begin{aligned}
s_1 : (q_1, p_1, q_2, p_2, t; \alpha_0, \alpha_1, \beta_0, \beta_1) \rightarrow &(-q_1, -(p_1 - 2q_1^2 - t), -q_2, -(p_2 - 2q_2^2 - t), t;\\
&\alpha_1, \alpha_0, \beta_1, \beta_0),\\
s_2 : (q_1, p_1, q_2, p_2, t; \alpha_0, \alpha_1, \beta_0, \beta_1) \rightarrow &(q_2, p_2, q_1, p_1, t; \beta_0, \beta_1, \alpha_0, \alpha_1).
\end{aligned}
$$

Theorem 1.5. *By the symplectic transformation* s_{II}

$$s_{II} : (q_1, p_1, q_2, p_2) \rightarrow (\frac{1}{(q_1 - q_2)}, -(q_1 - q_2)((q_1 - q_2)p_1 + \alpha_1), q_2, p_2 + p_1),$$

we can obtain the Hamiltonian system

$$\left\{ \begin{aligned} \frac{dq_1}{dt} = \frac{\partial H_{polynomial}}{\partial p_1} =& 2q_1^4 p_1 + 2\alpha_1 q_1^3 + p_2 q_1^2 + 2q_1 q_2 + 1, \\ \frac{dp_1}{dt} = -\frac{\partial H_{polynomial}}{\partial q_1} =& -4q_1^3 p_1^2 - 6\alpha_1 q_1^2 p_1 - 2(\alpha_1^2 + g4^2)q_1 \\ & -2q_1 p_1 p_2 - 2p_1 q_2 - \alpha_1 p_2, \\ \frac{dq_2}{dt} = \frac{\partial H_{polynomial}}{\partial p_2} =& -q_2^2 + p_2 - \frac{t}{2} + q_1(q_1 p_1 + \alpha_1), \\ \frac{dp_2}{dt} = -\frac{\partial H_{polynomial}}{\partial q_2} =& 2q_2 p_2 - 2q_1 p_1 - \alpha_1 + \beta_1 \end{aligned} \right. \tag{26}$$

with the Hamiltonian

(27)

$$\begin{aligned} H_{polynomial} =& q_1^4 p_1^2 + 2\alpha_1 q_1^3 p_1 + (\alpha_1^2 + g_4^2)q_1^2 + p_1 \\ & - q_2^2 p_2 + \frac{p_2^2}{2} + (\alpha_1 - \beta_1)q_2 - \frac{tp_2}{2} + q_1(q_1 p_1 p_2 + 2p_1 q_2 + \alpha_1 p_2). \end{aligned}$$

We remark that for this system we tried to seek its first integrals of polynomial type with respect to q_1, p_1, q_2, p_2. However, we can not find. Of course, the Hamiltonian $H_{polynomial}$ is not the first integral.

We also find some Bäcklund transformations of the system (26).

Theorem 1.6. *The system* (26) *becomes again a polynomial Hamiltonian system by each symplectic transformation*:

$$\begin{aligned} r_1 :& x_1 = (q_1 q_2 + 1)q_2, \ y_1 = \frac{p_1}{q_2^2}, \ z_1 = \frac{1}{q_2}, \\ & w_1 = -\left(q_2 p_2 - 2\left(q_1 q_2 + \frac{1}{2}\right)\left(\frac{p_1}{q_2}\right) - (\alpha_1 - \beta_1)\right) q_2, \\ r_2 :& x_2 = q_1, \ y_2 = p_1 - \frac{\alpha_0 - \alpha_1}{q_1} + \frac{2q_2^2 + t}{q_1^2} + \frac{4q_2}{q_1^3} + \frac{2}{q_1^4}, \\ & z_2 = q_2, \ w_2 = p_2 - 4q_2^2 - 2t - \frac{4q_2}{q_1} - \frac{2}{q_1^2}. \end{aligned} \tag{28}$$

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Megro-ku, 153-8914, Tokyo, Japan

E-mail address: `sasano@ms.u-tokyo.ac.jp`